

Research Studies Used for Vitamin D Deficiency Article

Abbas S, Chang-Claude J, Linseisen J. Plasma 25-hydroxyvitamin D and premenopausal breast cancer risk in a German case-control study. *Int J Cancer*. 2008.

Abbas S, Linseisen J, Slinger T, Kropp S, Mutschelknauss EJ, Flesch-Janys D, Chang-Claude J. Serum 25-hydroxyvitamin D and risk of post-menopausal breast cancer--results of a large case-control study. *Carcinogenesis*. 2008 ;29:93-9.

Bertone-Johnson, E.R. et al. Plasma 25-hydroxyvitamin D and 1, 25-dihydroxyvitamin D and risk of breast cancer. *Cancer Epidemiol. Biomarkers Prev*. 2005; 14: 1991-7.

Berwick M, Armstrong BK, Ben-Porat L, Fine J, Krickler A, Eberle C, Barnhill R. Sun exposure and mortality from melanoma. *J Natl Cancer Inst*. 2005;97:195-9.

EURODIAB. *Diabetes*. 1988 Aug;37(8):1113-9.

Feskanich D, et al. Plasma vitamin D metabolites and risk of colorectal cancer in women. *Cancer Epidemiol Biomarkers Prev* 2004;13:1502– 8.

Forman, JP et al. Plasma 25(OH)D and 4-year Multivariable Adjusted RR for Incident Hypertension. *Hypertension* 2007; 49:1063-9.

Freedman DM, Looker AC, Shih-Chen C, et al. Prospective study of serum vitamin D and cancer mortality in the United States. *J Natl Cancer Inst* 2007;99:1594-602.

Garland CF, et al. Vitamin D and prevention of breast cancer: Pooled analysis, *J Steroid Biochem Mol Biol*. 2007;103:708-11.

Giovannucci E, Liu Y, Rimm EB, Hollis BW, Fuchs CS, Stampfer MJ, Willett WC. Prospective study of predictors of vitamin D status and cancer incidence and mortality in men. *J Natl Cancer Inst*. 2006;98:451-9.

Giovannucci E, et al. Low 25-(OH)D levels associated with increased risk of MI in men. *Arch Intern Med* 2008;168:1174-80.

Goodwin PJ, et al. Vitamin D deficiency is common at breast cancer diagnosis and is associated with a significantly higher risk of distant recurrence and death in a prospective cohort study. American Society of Clinical Oncology Annual Meeting, Chicago, Illinois, May 30-June 3, 2008. Abstract number: 08-AB-31397-ASCOAM.

Gorham ED, et al. Optimal vitamin D status for colorectal cancer prevention: A Quantitative Meta Analysis. *Am J Prev Med* March 2007; 32:210-6.

Gorham ED, Mohr SB, Garland CF, Chaplin G, Garland FC. Do sunscreens increase risk of melanoma in populations residing at higher latitudes? *Ann Epidemiol*. 2007;17:956-63.

Gruber BM, Anuszevska EL. Influence of vitamin D3 metabolites on cell proliferation and cytotoxicity of adriamycin in human normal and neoplastic cells. *Toxicol In Vitro*. 2002;16:663-7.

Hathcock JN et al. Vitamin D intake and toxicity. *Am J Clin Nutr*. 2007;85:6-18.
**cites no toxicity below 30,000IU/day.

Heaney RP, Davies KM, Chen TC, Holick MF, Barger-Lux MJ. Human serum 25-hydroxycholecalciferol response to extended oral dosing with cholecalciferol. *Am J Clin Nutr*. 2003;77:204-10.

Hollis BW. Circulating 25-hydroxyvitamin D levels indicative of vitamin D sufficiency: implications for establishing a new effective dietary intake recommendation for vitamin D. *J Nutr*. 2005;135:317-22.

Hypertension 1997, 30:150-156. Shows epidemiology for hypertension by latitude.

HyppönenE, LääräE, ReunanenA, JärvelinMR, Virtanen SM. Intake of vitamin D and risk of type 1 diabetes: birth-cohort study. *Lancet*. 2001;358:1500-3.

Incidence Rates of Type I Diabetes in 51 Regions Young men, ages 0-18 by latitude, 1990-1994. World Health Organization.

Lappe JM, Travers-Gustafson D, Davies KM, Recker RR, Heaney RP. Vitamin D and calcium supplementation reduces cancer risk: results of a randomized trial. *Am J Clin Nutr*. 2007;85:1586-91.

Lowe LC, et al. Plasma 25-hydroxy vitamin D concentrations, vitamin D receptor genotype and breast cancer risk in a UK Caucasian population. *Eur J Cancer*. 2005;41:1164-9.

Martins D et al. Low 25-(OH)D Levels are Associated with the Metabolic Syndrome. *Arch Intern Med* 2007;167:1159-165.

Melamed M et al. Low 25-OHD Levels Associated with Higher Prevalence of Peripheral Arterial Disease: NHANES 2001–2004 (n= 4839) *Arterioscler Thromb Vasc Biol* 2008;28:1179-85.

Millen AE, Tucker MA, Hartge P, et al. Diet and melanoma in a case-control study. *Cancer Epidemiol Biomark Prev* 2004;13:1042-51.

Mohr SB, Gorham ED, Garland CF, Grant WB, Garland FC. Are low ultraviolet B and high animal protein intake associated with risk of renal cancer? *Int J Cancer*. 2006;119:2705-9.

Pelczynska M, Switalska M, Maciejewska M, Jaroszewicz I, Kutner A, Opolski A. Antiproliferative activity of vitamin D compounds in combination with cytostatics. *Anticancer Research* 2006;26:2701-5.

Plitz et al. Low vitamin D levels independently predict stroke. *Stroke* 2008;39:2611-13.

Stene LC, Joner G. Use of cod liver oil during the first year of life is associated with lower risk of childhood-onset type 1 diabetes: a large, population-based, case-control study. *Am J Clin Nutr.* 2003;78:1128-34.

Tangrea J, et al. Serum levels of vitamin D metabolites and the subsequent risk of colon and rectal cancer in Finnish men. *Cancer Causes Control* 1997;8:615–25.

Targher G. et al. Serum 25-(OH)D Levels in Type II diabetics: Correlation with carotid IMG. *Clin Endocrinol (Oxf)* 2006;65:593-7.

Twoogor SS, Lee IM, Buring JE, Rosner B, Hollis BW, Hankinson SE. Plasma 25-hydroxyvitamin D and 1,25-dihydroxyvitamin D and risk of incident ovarian cancer. *Cancer Epidemiol Biomarkers Prev.* 2007;16:783-8.

Wactawski-Wende J, et al. Calcium plus vitamin D supplementation and the risk of colorectal cancer. *New Engl J Med* 2006; 354:684-96.